

Summary of proposed changes to **Warnings** section of the Drug Facts panel with changes highlighted in red.

New/Revised Warnings	Previous Warnings
<p><b>Warnings</b>  <b>Allergy alert:</b> Ketoprofen may cause a severe allergic reaction which may include: ■ hives            ■ facial swelling ■ asthma (wheezing) ■ shock  <b>Stomach bleeding warning:</b> Taking more than recommended may cause stomach bleeding.  <b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ketoprofen or other pain relievers/fever reducers. Ketoprofen may cause stomach bleeding.</p>	<p><b>Warnings</b>  <b>Allergy alert:</b> Ketoprofen may cause a severe allergic reaction which may include: ■ hives            ■ facial swelling ■ asthma (wheezing) ■ shock  <b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ketoprofen or other pain relievers/fever reducers. Ketoprofen may cause stomach bleeding.</p>
<p><b>Do not use</b> if you have ever had an allergic reaction to any other pain reliever/fever reducer</p>	<p><b>Do not use</b> if you have ever had an allergic reaction to any other pain reliever/fever reducer</p>
<p><b>Ask a doctor before use if you have</b>            ■ problems or serious side effects from taking pain relievers or fever reducers            ■ stomach problems that last or come back, such as heartburn, upset stomach, or pain            ■ ulcers            ■ bleeding problems            ■ high blood pressure, heart or kidney disease, are taking a diuretic, or are over 65 years of age</p>	<p><b>Ask a doctor before use if you have</b> had problems or side effects with any pain reliever/fever reducer</p>
<p><b>Ask a doctor or pharmacist before use if you are</b>            ■ under a doctor's care for any serious medical condition            ■ taking any other product that contains ketoprofen or any other pain reliever/fever reducer            ■ taking a prescription drug for anticoagulation (blood thinning)            ■ taking any other drug</p>	<p><b>Ask a doctor or pharmacist before use if you are</b>            ■ under a doctor's care for any continuing medical condition            ■ taking other drugs on a regular basis            ■ taking any other product containing ketoprofen, or any other pain reliever/fever reducer</p>
<p><b>When using this product</b> take with food or milk if stomach upset occurs</p>	
<p><b>Stop use and ask a doctor if</b>            ■ an allergic reaction occurs. Seek medical help right away.            ■ pain get worse or lasts more than 10 days            ■ fever gets worse or lasts more than 3 days            ■ stomach pain or upset gets worse or lasts            ■ redness or swelling is present in the painful area            ■ any new symptoms appear</p>	<p><b>Stop use and ask a doctor if</b>            ■ an allergic reaction occurs. Seek medical help right away.            ■ fever gets worse or lasts more than 3 days            ■ pain gets worse or lasts more than 10 days            ■ stomach pain occurs with the use of this product            ■ the painful area is red or swollen            ■ any new or unexpected symptoms occur</p>